

A VARIANT OF SHANBHAG'S LEMMA ARISING OUT OF A MODIFIED
RAO-RUBBIN CONDIT. (U) PITTSBURGH UNIV PA CENTER FOR
MULTIVARIATE ANALYSIS A A ALZAID ET AL MAY 85

MULTIVARIATE ANALYSIS A A ALZAID ET AL. MAY 85

TR-85-22 AFOSR-TR-85-0558 F49620-85-C-0008 F/G 12/1

NL

52

END

PLATE 1

DYE

AFOSR-TR- 85 - 0558

AD-A158 608

A VARIANT OF SHANBHAG'S LEMMA ARISING
OUT OF A MODIFIED RAO-RUBIN CONDITION


by

Abdulhamid A. Alzaid, C. Radhakrishna Rao
and D. N. Shanbhag

King Saud University, Saudia Arabia,
University of Pittsburgh, U.S.A. and
University of Sheffield, U.K.

Center for Multivariate Analysis

University of Pittsburgh



DTIC
ELECTE
AUG 2 9 1985
S D
G

Approved for public release;
distribution unlimited.

DTIC FILE COPY

85 8 21 157

Unclassified

AD A 158 608

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER AFOSR-TR-85-0558	
4. PERFORMING ORGANIZATION REPORT NUMBER		6. NAME OF MONITORING ORGANIZATION AFOSR	
6a. NAME OF PERFORMING ORGANIZATION University of Pittsburgh	6b. OFFICE SYMBOL if applicable	7a. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING NUMBERING ORGANIZATION AFOSR	8b. OFFICE SYMBOL if applicable NM	9. PROGRAM ELEMENT, REPORT, and/or Contract Number F49620-85-C-0008	
9a. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448		10. SOURCE OF FUNDING NOS. PROGRAM ELEMENT NO. 61102F PROJECT NO. 2304 TASK NO. A5 WORK UNIT NO.	
11. TITLE (Include Security Classification) A Variant of Shanbhag's Lemma Arising Out of a Modified Rao-Rubbin Condition			
12. PERSONAL AUTHOR(S) Abdulhamid A. Alzaid, C. Radhakrishna Rao and D. N. Shanbhag			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Yr., Mo., Day) May 1985	15. PAGE COUNT 10

16. SUPPLEMENTARY NOTATION

17. DESCRIPTORS FIELD GROUP SUB GR XXXXXXXXXXXX	18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Damage models, Integrated Cauchy functional equation, Rao-Rubbin condition, Shanbhag's lemma
---	---

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

A modified Rao-Rubbin condition for damage models gives rise to a recurrence relation which is somewhat different from that considered by Shanbhag (1977). A complete solution to the new recurrence relation is obtained and its applications are indicated.

Additional figures in Random variables

Reproduced from
best available copy.

20. DISTRIBUTION STATEMENT (If the report is not available, indicate the reason)		21. DISTRIBUTION STATEMENT (If the report is not available, indicate the reason)	
Unclassified		Unclassified	
22. DISTRIBUTION STATEMENT (If the report is not available, indicate the reason)		23. DISTRIBUTION STATEMENT (If the report is not available, indicate the reason)	
Unclassified		Unclassified	

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSO)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for publication in the AD 199-12.
Distribution is unlimited.
MATTHEW J. KESLER
Chief, Technical Information Division

A VARIANT OF SHANBHAG'S LEMMA ARISING
OUT OF A MODIFIED RAO-RUBIN CONDITION

by

Abdulhamid A. Alzaid, C. Radhakrishna Rao
and D. N. Shanbhag

King Saud University, Saudia Arabia,
University of Pittsburgh, U.S.A. and
University of Sheffield, U.K.

May 1985

Technical Report No. 85-22

Center for Multivariate Analysis
515 Thackeray Hall
University of Pittsburgh
Pittsburgh, PA 15260

DTIC
ELECTE
AUG 29 1985
S D
G

*The work of this author was supported by the Air Force Office of Scientific Research under Contract #F49620-85-C-0008. The United States Government is authorized to reproduce and distribute reprints for governmental purposes not withstanding any copyright notation herein.

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

A VARIANT OF SHANBHAG'S LEMMA ARISING
OUT OF A MODIFIED RAO-RUBIN CONDITION

by

Abdulhamid A. Alzaid, C. Radhakrishna Rao
and D.N. Shanbhag

King Saud University, Saudi Arabia,
University of Pittsburgh, U.S.A., and
University of Sheffield, U.K.

Summary A modified Rao-Rubin condition for damage models gives rise to a recurrence relation which is somewhat different from that considered by Shanbhag (1977). A complete solution to the new recurrence relation is obtained and its applications are indicated.

Key Words: Damage models, Integrated Cauchy functional equation, Rao-Rubin condition, Shanbhag's lemma.

AMS Classification Index 60E05, 62E10

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A/	



1. INTRODUCTION

Let (X, Y) be a 2-vector of non-negative integer valued random variables such that $P(X \geq Y) = 1$ and the conditional distribution of Y given $X = n$ is binomial with index n and probability of success π . In such a case, the Rao-Rubin condition (see Rao and Rubin (1964))

$$P(Y = y) = P(Y = y | X = Y), \quad y = 0, 1, \dots \quad (1.1)$$

characterizes the distribution of X as Poisson. Shanbhag (1977) considered a more general convolution type conditional distribution of Y given X , in which case the condition (1.1) gives rise to the recurrence relation

$$v_m = \sum_{n=0}^{\infty} v_{m+n} w_n, \quad m = 0, 1, \dots \quad (1.2)$$

He obtained a complete solution of (1.2) in the form

$$v_n = v_0 b^n, \quad \sum_{n=0}^{\infty} b^n w_n = 1 \quad (1.3)$$

provided $v_n \neq 0$ for some $n \geq 1$ and $w_1 > 0$. Shanbhag's lemma proved a useful tool in solving a variety of characterization problems (see Rao and Shanbhag (1984) and the references cited in the paper).

A question has been raised by Srivastava and Singh (1975) as to whether the Rao-Rubin result on the Poisson distribution holds under the modified Rao-Rubin condition

$$P(Y = y) = P(Y = y | X - Y = k), \quad k > 0, \quad y = 0, 1, \dots \quad (1.4)$$

Patil and Taillie (1979) showed that the Rao-Rubin result may not hold in such a case, but two conditions of the type (1.4) for k and $k + k_1$ where $k_1 > 0$ provide a unique characterization of the distribution of X . Shanbhag and Taillie (1979)

(1979) extended the result of Patil and Taillie to a more general conditional distribution of Y given X . It is seen that a single condition of the type (1.4) leads to the recurrence relation

$$v_{m+k} = \sum_{n=0}^{\infty} v_{m+n} w_n, \quad m=0,1,\dots \quad (1.5)$$

where $k > 0$ is a fixed number. In such a case, the solution is not so simple. We obtain the complete solution to (1.5) and consider some applications.

The continuous analogues of the equations (1.2) and (1.5) have been considered by Lau and Rao (1982, 1984).

2. THE MAIN LEMMA

Let $\{(v_n, w_n): n=0,1,\dots\}$ be a sequence of vectors such that at least one $v_n \neq 0$ and $w_0 > 0$. Further let k be a positive integer such that the least common divisor of k and those n for which $w_n > 0$ be 1, and T be a non-negative matrix (in the sense of Senata (1973)) such that the corresponding state space is $\{0,1,2,\dots\}$ and the (i,j) -th element is

$$T_{ij} = \begin{cases} \delta_{ij} & \text{if } i,j=0,\dots,k-1 (\delta_{ii}=1 \text{ and } \delta_{ij}=0, i \neq j), \\ w_{j-i+k} & \text{if } i=k,k+1,\dots \text{ and } j \geq i-k, \\ 0 & \text{otherwise.} \end{cases}$$

Define by $\{f_{i,r}: i \geq k, r=0,\dots,k-1\}$ the sequence of absorption measures corresponding to T and by $T_{ij}^{(n)}$, the n -step transition measure corresponding to the transition $i \rightarrow j$. Let

$$D = \{b: b > 0, b^k = \sum_{n=0}^{\infty} b^n w_n\}$$

LEMMA The sequence $\{(v_n, w_n): n = 0, 1, \dots\}$ as defined above satisfies the recurrence equations

$$v_{m+k} = \sum_{n=0}^{\infty} w_n v_{n+m}, \quad m = 0, 1, \dots \quad (2.1)$$

iff D is non-empty, $f_{m,0}, \dots, f_{m,k-1}$ are finite, and one of the following hold.

(i) D has only one point, $\sum_{n=0}^{\infty} (n-k)b^n w_n < 0$ for $b \in D$, and

$$v_m = f_{m,0}v_0 + \dots + f_{m,k-1}v_{k-1}, \quad m = k, k+1, \dots$$

(ii) D has only one point, $\sum_{n=0}^{\infty} (n-k)b^n w_n = 0$ for $b \in D$ and for some $c \geq 0$

$$v_m = f_{m,0}(v_0 - c \cdot 0 \cdot b^0) + \dots + f_{m,k-1}(v_{k-1} - c(k-1)b^{k-1}) + cmb^m,$$

$$m = k, k+1, \dots,$$

with $b \in D$.

(iii) D contains two points and for some $c \geq 0$

$$v_m = f_{m,0}(v_0 - cb^0) + \dots + f_{m,k-1}(v_{k-1} - cb^{k-1}) + cb^m, \quad m = k, k+1, \dots,$$

with b as the larger of the two members of D .

Proof The 'if' part of the theorem follows by a straight-forward verification.

We shall now establish the 'only if' part of the theorem. Let $V = (v_0, v_1, \dots)'$.

Clearly we have $V = TV$ and hence

$$V = T^n V, \quad n \geq 1. \quad (2.2)$$

Observe that $\{T_{ij}^{(n)}\}$ is an increasing sequence in n for each $i \geq k$ and $j = 0, 1, \dots, k-1$.

Consequently, $\lim_{n \rightarrow \infty} T_{ij}^{(n)}$ exists for each $i \geq k$ and $j = 0, 1, \dots, k-1$. These are given

by $f_{i,j}$ defined earlier. Then (2.2) implies that

$$v_m = f_{m,0} v_0 + \dots + f_{m,k-1} v_{k-1} + \lim_{n \rightarrow \infty} \sum_{j=k}^{\infty} T_{mj}^{(n)} v_j, \quad m = k, k+1, \dots \quad (2.3)$$

Since at least one $v_n \neq 0$, it is clear that there exists an $n_0 \geq k$ such that $v_{n_0} > 0$. Given any $j \in \{0, 1, \dots, k-1\}$, clearly (2.3) is valid with $\{v_m\}$ replaced by $\{v_m^{(j)}\}$ where

$$v_m^{(j)} = v_{m+n_0-j}, \quad m \geq 0.$$

Consequently, it follows in view of (2.3) that $f_{m,j} < \infty$ for each $m \geq k$ and $j = 0, 1, \dots, k-1$. Replacing v_m in (2.3) by v_{m+r-k} , we get for $m = k$

$$v_r = f_{k,0} v_{r-k} + \dots + f_{k,k-1} v_{r-1} + \lim_{n \rightarrow \infty} \sum_{j=k}^{\infty} T_{kj}^{(n)} v_{j+r-k}, \quad r \geq k. \quad (2.4)$$

Define

$$\xi_m^{(n)} = \sum_{j=k}^{\infty} T_{kj}^{(n)} v_{j+m-k}, \quad m = 0, 1, 2, \dots$$

and $\xi_m = \lim_{n \rightarrow \infty} \xi_m^{(n)}$. In view of the dominated convergence theorem, the existence of $\lim_{n \rightarrow \infty} \xi_m^{(n)}$ for $m = 0, 1, \dots, k-1$ follows inductively from the fact that the sequence $\{\xi_m^{(n)}\}_{m=0}^{\infty}$ is a non-negative real sequence satisfying (2.1) for each n .

By considering t to be the minimum of the states in the passage from k to j , we get

$$\begin{aligned} \xi_m^{(n)} &= \sum_{j=k}^{\infty} T_{kj}^{(n)} v_{j+m-k} \\ &= \sum_{l=0}^{\infty} w_{l+k} \sum_{r=0}^{n-1} \sum_{t=k}^{l+k} t_{f_{l+k,t}}^{(r)} \sum_{j=t}^{\infty} T_{k,j-t+k}^{(n-r-1)} v_{j+m-k} \\ &= \sum_{l=0}^{\infty} w_{l+k} \sum_{r=0}^{n-1} \sum_{t=k}^{l+k} t_{f_{l+k,t}}^{(r)} \xi_{m+t-k}^{(n-r-1)}, \end{aligned} \quad (2.5)$$

where $t_{f_{l+k,t}}^{(r)}$ is the measure of the set that the first passage from $l+k$ to t avoiding states lower than t takes place at time r . The term following w_{l+k} under the summation on the right hand side of (2.5) is bounded by v_{l+m} .

Consequently, the dominated convergence theorem implies that

$$\begin{aligned}\xi_m &= \sum_{l=0}^{\infty} w_{l+k} \sum_{r=0}^{\infty} \sum_{t=k}^{l+k} t_{f_{l+k,t}}^{(r)} \xi_{m+t-k} \\ &= \sum_{t=k}^{\infty} \xi_{m+t-k} \sum_{l=t-k}^{\infty} w_{l+k} \sum_{r=0}^{\infty} t_{f_{l+k,t}}^{(r)}.\end{aligned}$$

From Shanbhag's lemma, we can then conclude that

$$\xi_m = \alpha b^m, \quad m = 0, 1, 2, \dots,$$

for some non-negative α and b . (For the case $w_l = 0$ for all $l \geq k+1$, the result is trivial; otherwise this is a corollary of Shanbhag's lemma.) Using (2.4), it follows that

$$v_m = f_{k,0} v_{m-k} + \dots + f_{k,k-1} v_{m-1} + \alpha b^m, \quad m \geq k. \quad (2.6)$$

If $\theta > 0$ such that

$$\theta^k = f_{k,0} \theta^0 + \dots + f_{k,k-1} \theta^{k-1}, \quad (2.7)$$

then clearly by induction (in view of the fact that $f_{k,0}, \dots, f_{k,k-1}$ are respectively the first passage measures for the transition $m \rightarrow m-k+j$ avoiding the states lower than m until then for $j = 0, 1, \dots, k-1$).

$$\begin{aligned}\theta^m &= f_{k,0} \theta^{m-k} + \dots + f_{k,k-1} \theta^{m-1} \\ &\vdots \\ &= f_{m,0} \theta^0 + \dots + f_{m,k-1} \theta^{k-1}, \quad m = k, k+1, \dots,\end{aligned}$$

which implies $\theta \in D$. Thus D is non-empty. If $\alpha \neq 0$ and $b \neq 0$, then $b \in D$. If D is a singleton and $\alpha, b \neq 0$, then b satisfies (2.7) and there exists a $c > 0$ such that

$$ckb^k = c \cdot 0 f_{k,0} b^0 + \dots + c(k-1) f_{k,k-1} b^{k-1} + \alpha b^k.$$

Consequently (2.6) implies that

$$\begin{aligned} v_m - cmb^m &= f_{k,0}(v_{m-k} - c(m-k)b^{m-k}) + \dots + f_{k,k-1}(v_{m-1} - c(m-1)b^{m-1}) \\ &\vdots \\ &= f_{m,0}(v_0 - c \cdot 0 \cdot b^0) + \dots + f_{m,k-1}(v_{k-1} - c(k-1)b^{k-1}), \quad m \geq k, \end{aligned}$$

which, in turn, implies that $\{cmb^m\}$ satisfies (2.1) and hence in view of the fact that $b \in D$ yields that $\sum_{n=0}^{\infty} (n-k)b^n w_n = 0$. On the other hand, if D is a doubleton and $\alpha, b \neq 0$, then essentially an argument the same as the above implies that b cannot satisfy (2.7) since $\sum_{n=0}^{\infty} (n-k)b^n w_n = 0$ implies D to be a singleton. The b in question is a point of D and hence we have

$$b^k > f_{k,0} b^0 + \dots + f_{k,k-1} b^{k-1},$$

which implies that there exists a $c \geq 0$ such that

$$cb^k = c \cdot b^0 f_{k,0} + \dots + cb^{k-1} f_{k,k-1} + \alpha b^k$$

or

$$\begin{aligned} v_m - cb^m &= f_{k,0}(v_{m-k} - cb^{m-k}) + \dots + f_{k,k-1}(v_{m-1} - cb^{m-1}) \\ &= f_{m,0}(v_0 - cb^0) + \dots + f_{m,k-1}(v_{k-1} - cb^{k-1}), \quad m \geq k. \end{aligned}$$

The result is now obvious. [Note that if D is a singleton and $b \in D$, then it is impossible that $\sum_{n=0}^{\infty} (n-k)b^n w_n = 0$.]

Recently Lau and Rao (1984) considered a continuous analogue of (2.1), which is a modified version of the integrated Cauchy functional equation discussed in Lau and Rao (1982), and obtained a general solution using a random walk approach.

Corollary If the sequence (2.1) is such that

$$\lambda v_{n+k} = v_{n+k_1}, \quad n=0,1,\dots,k,$$

for some integer $k_1 > k$ and some real λ , then

$$v_n = v_0 \lambda^{n/(k_1-k)}, \quad n=0,1,\dots$$

Proof The result of the main lemma (in particular, the relation (2.6)) implies that

$$v_{n+k_1-k} = \lambda v_n \quad \text{for all } n \geq 0 \quad (2.8)$$

with $\lambda > 0$. Reducing the system of equations in (2.6) to the one in Shanbhag's lemma by appropriate substitutions via (2.8) we arrive at the required result.

Note 1. It is also possible to prove the result of the corollary by using Perron-Frobenius theorem given in Seneta (1973, pp. 1-2). For details, the reader is referred to Alzaid (1983).

Note 2. Let $Y_{1,n} \leq \dots \leq Y_{n,n}$ denote ordered observations in a sample of size n from a non-degenerate distribution F concentrated on non-negative integers. Arnold (1980) raised the question whether the independence of the r.v. $Y_{2,n} - Y_{1,n}$ and the event $\{Y_{1,n} = m\}$ for a fixed $m > 1$ implies the distribution F to be geometric. It is easily seen that this property leads to a recurrence relation of the type (2.1). Hence an application of the main lemma shows that the distribution is not necessarily geometric. However, it can be shown that if the stated independence holds for two fixed values of m , then under some mild conditions, F is indeed geometric.

Note 3. In view of the result of our main lemma, it is seen that Theorem 4 of Krishnaji (1974) is not correct. This also follows from the counter example given by Patil and Taillie (1979). The error in Krishnaji's argument appears in the last sentence of the proof in which it is claimed that since $X = \Lambda \exp\{\Lambda(\theta-1)\}$ is degenerate, Λ has to be degenerate. We may however point out here that Krishnaji's theorem with the portion " $G(t)$ is non-negative for all real t " in it replaced by " $G(t)$ is infinitely divisible" is valid.

Note 4. The result of our main lemma could also be applied to obtain certain conclusions concerning stationary queue length distributions in $GI/E_k/1$ queueing systems and also in their modifications when the arrivals are in batches of random size. In particular, the result implies that the queue length distribution corresponding to such a system exists if and only if the relative traffic intensity of the system is less than 1 and also that the stationary queue length distribution for the system $GI/M/1$ is geometric.

Note 5. Let (X, Y) be a random vector variable with non-negative integer valued components such that the conditional distribution of Y given $X=x$ is given by

$$S(y|x) = \frac{a_y b_{x-y}}{c_x}, \quad y = 0, 1, \dots, x$$

for almost all x , where $\{c_n\}$ is the convolution of $\{a_n\}$ and $\{b_n\}$, $a_n > 0$ for all n , $b_0 > 1$ and $b_n \geq 0$, $n \geq 1$ with the least common divisor of n for which $b_n > 0$ equal to 1. Then, according to the corollary to the main lemma, the conditions

$$P(Y=y) = P(Y=y | X-Y=k_0), \quad y = 0, 1, \dots,$$

$$P(Y=y | X-Y=k_0) = P(Y=y | X-Y=k_0+k_1), \quad y = 0, 1, \dots, k_0,$$

for some $0 \leq k_0 < k_0 + k_1 < \infty$ with $P(X-Y=k_0) > 0$ and $P(X-Y=k_0+k_1) > 0$ is equivalent to the assertion that Y and $X-Y$ are independent. This implies that

$$P(X=x) \propto c_x \theta^x, \quad x = 0, 1, 2, \dots$$

for some $\theta > 0$.

The above result extends an unpublished characterization given by Shanghag and Taillie (1979) of a distribution $\{g_x\}$ of the type

$$g_x \propto c_x \theta^x, \quad x = 0, 1, \dots \quad (2.9)$$

with θ as some positive constant, based on two modified Rao-Rubin conditions. Our lemma proves that there exist infinitely many distributions other than that of the form (3.9) for which one modified Rao-Rubin condition holds and implicitly identifies the class of such distributions.

3. REFERENCES

- [1] Alzaid, A.A. (1983). Some Contributions to Characterization Theory. Ph.D. Thesis, University of Sheffield.
- [2] Arnold, B.G. (1980). Two characterizations of the geometric distribution. J. Appl. Prob. 17, 570-573.
- [3] Krishnaji, N. (1974). Characterization of some discrete distributions based on a damage model. Sankhya A, 36, 204-213.
- [4] Lau, Ka-Sing and Rao, C.R. (1982). Integrated Cauchy functional equation and characterizations of the exponential law. Sankhya A, 44, 72-90.
- [5] Lau, Ka-Sing and Rao, C.R. (1984). On the integral equation $f(x) = \int_R f(x+y) d\mu(y)$, $y \geq 0$. Tech. Rept. 84-09, University of Pittsburgh.
- [6] Patil, G.P. and Taillie, C. (1979). On a variation of the Rao-Rubin theorem. Sankhya A, 41, 129-132.
- [7] Rao, C.R. and Shanbhag, D.N. (1984). Recent results on characterization of probability distributions: a unified approach through extensions of Deny's theorem. Tech. Rept. 84-39, University of Pittsburgh.

- [8] Rao, C.R. and Rubin, H. (1964). On a characterization of the Poisson distribution. Sankhya A, 26, 295-298.
- [9] Seneta, E. (1973). Non-negative Matrices, George Allen and Unwin Ltd., London.
- [10] Shanbhag, D.N. (1977). An extension of Rao-Rubin characterization of the Poisson distribution. J. Appl. Prob. 14, 640-646.
- [11] Shanbhag, D.N. and Taillie, C. (1979). An extension of the Patil-Taillie characterization of the Poisson distribution (unpublished).
- [12] Srivastava, R.C. and Singh, J. (1975). On characterizations of the binomial and Poisson distributions based on a damage model. In Statistical Distributions in Scientific Work (Eds. Patil, Kotz and Ord), D. Reidel Publishing Company, pp).

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 85-22	2. GOVT ACCESSION NO. AD-A158 608	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Variant of Shanbhag's Lemma Arising Out of a Modified Rao-Rubbin Condition		5. TYPE OF REPORT & PERIOD COVERED Technical - May 1985
		6. PERFORMING ORG. REPORT NUMBER 85-22
7. AUTHOR(s) Abdulhamid A. Alzaid, C. Radhakrishna Rao and D. N. Shanbhag		8. CONTRACT OR GRANT NUMBER(s) F49620-85-C-0008
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis 515 Thackeray Hall University of Pittsburgh, Pgh, PA 15260		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Department of the Air Force Bolling Air Force Base, DC 20332		12. REPORT DATE May 1985
		13. NUMBER OF PAGES 10
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		16. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Damage models, Integrated Cauchy functional equation, Rao-Rubin condition, Shanbhag's lemma		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A modified Rao-Rubbin condition for damage models gives rise to a recurrence relation which is somewhat different from that considered by Shanbhag (1977). A complete solution to the new recurrence relation is obtained and its applications are indicated.		

DD FORM 1 JAN 73 1473

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

END

END

FILMED

10-85

DTIC